

Table S2. Aetiological diagnoses of the SSTIs following allo-HSCT.

Type	Organism	No. (%)
Bacteria		18 (32.7%)
	<i>Staphylococcus aureus</i>	2 (3.6%)
	<i>Staphylococcus epidermidis</i>	1, (1.8%)
	<i>Staphylococcus haemolyticus</i>	2, (3.6%)
	<i>Staphylococcus epidermidis</i>	2, (3.6%)
	<i>Enterococcus faecium</i>	3, (5.5%)
	<i>Escherichia coli</i>	6, (10.9%)
	<i>Klebsiella pneumoniae</i>	1, (1.8%)
	<i>Corynebacterium diphtheriae</i>	1, (1.8%)
	<i>Acinetobacter baumannii</i>	1, (1.8%)
Fungi		12, (21.8%)
	undefined	6, (10.9%)
	mycete, undefined	1, (1.8%)
	Mucor	1, (1.8%)
	<i>Aspergillus fumigatus</i>	2, (3.6%)
	<i>Candida Albicans</i>	1, (1.8%)
	<i>Candida Glabrata</i>	1, (1.8%)
Viruses		25, (45.5%)
	HSV-1	9, (16.4%)
	HSV-2	5, (9.1%)
	VZV	10, (18.2%)
	HPV	1, (1.8%)

Notes: HSV, herpes simplex virus; VZV, varicella-zoster virus; HPV, Human Papilloma Virus